



SEPSIS ALLIANCE®

Suspect Sepsis. Save Lives.™

Life After Sepsis Fact Sheet

Definition: Sepsis is the body's overwhelming and life-threatening response to infection, which can lead to tissue damage, organ failure, and death.

Who it Hurts: While sepsis is an equal-opportunity killer, impacting the sick, the well, and people of all ages, some groups are more likely to be affected. These include very young children, older adults, and those with a weakened immune system.

Prevention: The risk of sepsis can be reduced by preventing or quickly identifying and managing infections. This includes practicing good hygiene, staying current with vaccinations, and seeking treatment when infections are suspected.

Treatment: Sepsis is a medical emergency that requires urgent attention and rapid treatment for survival. Sepsis can be treated and, in many instances, lives are saved by using existing and proven protocols.

Recovery: Many individuals fully recover from sepsis, while many others are left with long-lasting effects, such as amputations or organ dysfunction, like kidney failure. Other after-effects of sepsis are less obvious, such as memory loss, anxiety, or depression.

Symptoms: When it comes to sepsis, remember **It's About TIME™**:

T – Temperature - higher or lower than normal

I - Infection – may have signs or symptoms of infection

M – Mental Decline - confused, sleepy, difficult to rouse

E – Extremely Ill – “I feel like I might die,” severe pain or discomfort

If you **suspect sepsis** (observe a combination of these symptoms), see your medical professional immediately, CALL 911, or go to a hospital with an advocate and say, “**I AM CONCERNED ABOUT SEPSIS.**”

Post-sepsis syndrome: Up to 50% of sepsis survivors are left with physical and/or psychological long-term effects, a condition known as post-sepsis syndrome.^{1,2,3} These effects include:

- Insomnia, difficulty getting to sleep or staying asleep
- Nightmares, vivid hallucinations and panic attacks
- Disabling muscle and joint pains
- Extreme fatigue
- Poor concentration
- Decreased mental (cognitive) functioning
- Loss of self-esteem and self-belief

Life After Sepsis in Adults

- About one-third of all patients and more than 40% of older patients have another hospitalization within three months of the initial sepsis, most commonly due to a repeat episode of sepsis or another infection.^{1,2}
- One-half to two-thirds of all hospital readmissions after sepsis are infection-related.⁴
- The higher risk of infection following sepsis results from suppression or weakening of the immune system in the first few weeks and months following the initial bout of sepsis.^{1,4}
- In addition to infection, other common causes of rehospitalization after sepsis are heart failure, kidney failure, and pulmonary aspiration (inhaling food into the lungs).²
- Almost 60% of sepsis survivors experience worsened cognitive (mental) and/or physical function.³
- Older sepsis patients experience on average 1 to 2 new limitations on activities of daily living (e.g. bathing, dressing, managing money) after hospitalization.³
- One-sixth of sepsis survivors experience difficulties with memory, concentration, and decision making.²
- Older severe sepsis survivors are at higher risk for long-term cognitive impairment and physical problems than others their age who were treated for other illnesses.^{3,5}
- Many sepsis survivors report symptoms of post-traumatic stress disorder.⁶
- Patients who have had sepsis are more likely to develop symptoms of post-traumatic stress disorder than other ICU survivors.⁷
- Younger adult sepsis survivors (under 45 years old) are at a higher risk of death during the first 2 years after hospital discharge than patients hospitalized for infection without sepsis. These sepsis survivors tend to have more chronic (co-occurring) conditions (comorbidities) that may contribute to this increased risk.⁸
- Patients recovering from sepsis are at increased risk of stroke and heart attack (myocardial infarction) in the first 4 weeks after hospital discharge. Younger patients (aged 20 to 45) are at a higher risk for these events compared to older patients (those over age 75).⁹
- Caregivers to patients who have survived critical illness are at risk for poor mental health outcomes. 67% of caregivers (family members or friends) report high levels of depressive symptoms.¹⁰

Life After Sepsis in Children

- Nearly half (47%) of child sepsis survivors need to be readmitted to the hospital at least once for additional care after surviving sepsis.¹¹
- Among pediatric sepsis survivors, almost one third (31%) are discharged from the hospital with some disability, including cognitive or physical impairments, skin graft, amputation, or hearing loss.¹²
- Almost 1 in 6 pediatric severe sepsis survivors are discharged from the hospital with more disabilities than when they were admitted.¹³
- Survival from sepsis can be very challenging with many children requiring amputations. Many more experience a decrease in cognitive and physical function, with 34 percent of pediatric sepsis survivors (more than 1 in 3) showing a decline in their functional status at 28 days after hospital discharge.¹⁴

- Some research points to problems in cognitive function (mental work) after critical illness in children. In one small study, children who had spent time in the pediatric ICU for critical illness showed deficiencies in tests of memory-related tasks compared to healthy children. Survivors of critical illness in this study also had declines in their academic performance as evaluated by teachers when compared to healthy children. In another study, 44% of children who had survived septic shock were found to have problems with cognitive function when compared to healthy children.^{15,16}
- Pediatric severe sepsis survivors who have spent time in the pediatric ICU experience post-traumatic stress disorder more often than children discharged from the general hospital ward.¹³
- Critical illness in children takes a toll on family and caregivers. Higher rates of post-traumatic stress disorder and symptoms of deteriorating physical and mental health such as headaches, fatigue, and anxiety are reported for parents of survivors of pediatric critical illness.¹³

To find out more please visit [Sepsis.org](https://www.sepsis.org)

Sources:

1. Prescott, H and Angus, D, *JAMA*. 2018;319(1):62-75.
<https://jamanetwork.com/journals/jama/article-abstract/2667727?redirect=true>
2. Prescott, H and Angus, D., *JAMA*. 2018;319(1):91. <https://jamanetwork.com/journals/jama/fullarticle/2667724>
3. Iwashyna, TJ et al., *JAMA Network*. 2010;304(16):8. <https://jamanetwork.com/journals/jama/fullarticle/186769>
4. Prescott HC and Costa DK.. *Crit Care Clin*. 2018;34(1):175-188.
[http://www.criticalcare.theclinics.com/article/S0749-0704\(17\)30078-7/abstract](http://www.criticalcare.theclinics.com/article/S0749-0704(17)30078-7/abstract)
5. <http://www.uofmhealth.org/news/1782sepsis-survivors-more-than-three-times-as>
6. Boer KR et al. *Intensive Care Med*. 2008;34(4):664-674.
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2271079/>
7. https://www.hopkinsmedicine.org/news/media/releases/ptsd_symptoms_common_among_icu_survivors
8. Abu-Kaf H, et al., *J. Crit Care Med*. 2018;46(8):1269-1275. <https://www.ncbi.nlm.nih.gov/pubmed/29742586>
9. Lai C-C, et al., *Can Med Assoc J*. 2018;190:E1062-1069.
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6131076/>
10. Cameron JI, et al., *N Engl J Med*. 2016;374(19):11. <https://www.nejm.org/doi/full/10.1056/NEJMoa1511160>
11. Czaja, AS et al. *Pediatrics*. 2009;123(3):849-857.
<http://pediatrics.aappublications.org/content/123/3/849.short>
12. Boeddha NP, et al., *Crit Care*. 2018;22(1):143. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5984383/>
13. Syngal P and Giuliano JS., *Healthcare (Basel)*. 2018;6(3).
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6164000/>
14. Farris RW, et al., *Pediatr Crit Care Med*. 2013;14(9):835-842.
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4080839/>
15. Als LC, et al., *Crit Care Med*. 2013;41(4):1094-1103. <https://www.ncbi.nlm.nih.gov/pubmed/23385103>
16. Bronner MB, et al., *Pediatr Crit Care Med*. 2009;10(6):636-642.
<https://www.ncbi.nlm.nih.gov/pubmed/19581821>